High school students who used tobacco by selected demographic characteristics

Category	Percent (95% CI)								
	Current Cigarette Use*				Smokeless Tobacco Use**				Current Cigar Use***
	1993	1995	1997	1999	1993	1995	1997	1999	1999
Total	30.5 (28.6, 32.4)	34.8 (32.6, 37.0)	36.4 (34.1, 38.7)	34.8 (32.3, 37.3)	11.5 (9.9, 13.1)	11.4 (9.7, 13.1)	9.3 (7.1, 11.5)	7.8 (5.6, 10.0)	17.7 (16.0, 19.4)
Race/ethnicity									
Non-Hispanic white	33.7 (31.5, 35.9)	38.3 (35.7, 40.9)	39.7 (37.3, 42.1)	38.6 (35.4, 41.8)	14.6 (12.8, 16.4)	14.5 (12.8, 16.2)	12.2 (9.7, 14.7)	10.4 (7.0, 13.8)	18.8 (16.4, 21.2)
Non-Hispanic black	15.4 (12.9, 17.9)	19.2 (16.2, 22.2)	22.7 (18.9, 26.5)	19.7 (15.6, 23.8)	2.6 (1.4, 3.8)	2.2 (1.2, 3.2)	2.2 (1.1, 3.3)	1.3 (0.2, 2.4)	13.7 (11.3, 16.1)
Hispanic	28.7 (25.5, 31.9)	34.0 (28.8, 39.2)	34.0 (31.3, 36.7)	32.7 (28.9, 36.5)	4.9 (3.6, 6.2)	4.4 (2.6, 6.2)	5.1 (2.8, 7.4)	3.9 (2.3, 5.5)	16.7 (14.2, 19.2)
Grade									
9th	27.8 (25.5, 30.1)	31.2 (29.5, 32.9)	33.4 (28.3, 38.5)	27.6 (23.6, 31.6)	10.5 (8.4, 12.6)	11.2 (9.5, 12.9)	9.7 (7.0, 12.4)	6.8 (3.6, 10.0)	13.7 (10.4, 17.0)
10th	28.0 (24.7, 31.3)	33.1 (29.3, 36.9)	35.3 (31.2, 39.4)	34.7 (32.2, 37.2)	11.2 (9.0, 13.4)	9.6 (7.4, 11.8)	6.8 (5.1, 8.5)	7.1 (4.6, 9.6)	17.8 (15.2, 20.4)
11th	31.1 (27.9, 34.3)	35.8 (32.2, 39.4)	36.6 (33.0, 40.2)	36.0 (33.0, 39.0)	11.8 (9.6, 14.0)	13.0 (10.3, 15.7)	10.0 (7.5, 12.5)	8.4 (6.0, 10.8)	18.2 (16.6, 19.8)
12th	34.5 (30.7, 38.3)	38.2 (34.7, 41.7)	39.6 (34.7, 44.5)	42.8 (37.3, 48.3)	12.1 (10.4, 13.8)	11.2 (8.4, 14.0)	10.5 (6.9, 14.1)	8.9 (5.4, 12.4)	22.0 (18.4, 25.6)
Sex									
Male	29.8 (27.5, 32.1)	35.4 (33.0, 37.8)	37.7 (35.0, 40.4)	34.7 (31.7, 37.7)	20.4 (17.4, 23.1)	19.7 (17.2, 22.2)	15.8 (12.1, 19.5)	14.2 (10.4, 18.0)	25.4 (23.1, 27.7)
Female	31.2 (29.1, 33.3)	34.3 (31.2, 37.4)	34.7 (31.9, 37.5)	34.9 (32.3, 37.5)	2.0 (1.4, 2.6)	2.4 (1.1, 3.7)	1.5 (0.8, 2.2)	1.3 (0.8, 1.8)	9.9 (7.7, 12.1)

*Smoked cigarettes on ≥1 of the 30 days preceding the survey.
**Used chewing tobacco or snuff during the 30 days preceding the survey.
***Smoked cigars on ≥1 of the 30 days preceding the survey.
Source: MMWR Surveillance Summaries for the 1993, 1995, 1997, and 1999 Youth Risk Behavior Surveillance System, Centers for Disease Control and Prevention.