

Incidence rates for oral and pharyngeal cancers by stage and selected demographic characteristics

SEER Historic Stage	Race		
	White	Black	All Races
	Rate per 100,000* (95% CI)		
In Situ	0.36 (0.33, 0.39)	0.23 (0.17, 0.31)	0.34 (0.31, 0.36)
Localized	3.78 (3.69, 3.87)	2.56 (2.33, 2.80)	3.56 (3.48, 3.64)
Regional	4.26 (4.16, 4.35)	6.72 (6.34, 7.10)	4.50 (4.41, 4.59)
Distant	0.70 (0.66, 0.74)	1.51 (1.34, 1.70)	0.77 (0.73, 0.81)
Unstaged	1.04 (1.00, 1.09)	1.45 (1.29, 1.63)	1.11 (1.07, 1.16)
All Stages	10.13 (9.99, 10.28)	12.47 (11.96, 12.99)	10.28 (10.14, 10.41)
SEER Historic Stage	Gender		
	Male		Female
	Rate per 100,000* (95% CI)		
In Situ	0.47 (0.43, 0.52)		0.22 (0.20, 0.25)
Localized	5.11 (4.97, 5.25)		2.31 (2.23, 2.40)
Regional	6.82 (6.66, 6.99)		2.51 (2.42, 2.60)
Distant	1.19 (1.12, 1.26)		0.41 (0.37, 0.45)
Unstaged	1.74 (1.66, 1.82)		0.57 (0.53, 0.61)
All Stages	15.34 (15.09, 15.58)		6.02 (5.88, 6.17)

* The rates are age-adjusted to the 1970 U.S. standard population.

Data source: Surveillance, Epidemiology, and End Results (SEER) 11 Registries Public-Use, August 1999 Submission (1992-1997).